

Study on a Variant of Newton's Method with Accelerated Third-Order Convergence

KAPIL, Former Students, Department of Mathematics, Kurukshetra University, Kurukshetra, Haryana (India),
 Email: kapilkaswan409@gmail.com

Abstract:

In the given method, we suggest an improvement to the iteration of Newton's method. Derivation of Newton's method involves an indefinite integral of the derivative of the function, and the relevant area is approximated by a rectangle. In the proposed scheme, we approximate this indefinite integral by a trapezoid instead of a rectangle, thereby reducing the error in the approximation. It is shown that the order of convergence of the new method is three, and computed results support this theory. Even though we have shown that the order of convergence is three, in several cases, computational order of convergence is even higher. For most of the functions we tested, the order of convergence in Newton's method was less than two and for our method, it was always close to three.

Keywords: Variant, Newton Method, Accelerated

Introduction

In this study, authors have suggested an improvement over the Newton's method at the expense of one additional first derivative evaluation. Derivation of Newton's method involves an indefinite integral of the derivative of the function, and the relevant area is approximated by a rectangle. Here authors have approximated this indefinite integral by a trapezoid instead of a rectangle, and the resulted method has third-order convergence, *i.e.*, the method approximately triples the number of significant digits after some iterations. Computed results overwhelmingly support this theory, and computational order of convergence was even more than three for certain functions.

It is important to understand how Newton's method is constructed. At each iterative step construct a local model of the function $f(x)$ at the point x_n and solve for the root x_{n+1} of the local model. In Newton's method, shown in figure 1.1, the local linear model is the tangent drawn to the function $f(x)$ at the contact point x_n as:

$$M_n(x) = f(x_n) + f'(x_n)(x - x_n). \quad \dots(1.1)$$

Dennis [1983] interpreted local linear model in another way. From Newton's theorem,

$$f(x) = f(x_n) + \int_{x_n}^x f'(\lambda) d\lambda. \quad \dots(1.2)$$




Figure 1.1 : Newton's iterative step.




Figure 1.2 : Approximating the area by the rectangle ABCD. E




Figure 1.3 : Approximating the area by the trapezoid ABCD.

Dennis [1983] replaced the indefinite integral by the rectangle ABCD as shown in figure 1.2, *i.e.*,

$$\int_{x_n}^x f'(\lambda) d\lambda \approx f'(x_n)(x - x_n), \quad \dots(1.3)$$

which results in the model given in equation (1.1). In this the area DCE is ignored.

A Variant of Newton's Method

In this paper the authors have approximated the indefinite integral involved in equation (1.2) by the trapezium ABED as shown in figure 1.3, *i.e.*,

$$\int_{x_n}^x f'(\lambda) d\lambda \approx \left(\frac{1}{2}\right)(x - x_n)[f'(x_n) + f'(x)] \quad \dots(1.4)$$

Thus, the local modal is given by

$$M_n(x) = f(x_n) + \left(\frac{1}{2}\right)(x - x_n)[f'(x_n) + f'(x)]. \quad \dots(1.5)$$

It is found that not only the model (equation (1.5)) and its derivative agree with the function $f(x)$ and its derivative $f'(x)$, respectively, but the second derivative of the model and the second derivative of the function are also agreeing at the current iterate $x = x_n$ (Fernando and Weerakoon [1997]). Even though the model for Newton's method matches with the values of the slope $f'(x_n)$ of the function, it does not match with its curvature in terms of $f''(x_n)$.

The next iterative point as the root of the local model (equation (1.5)) is

$$\begin{aligned} M_n(x_{n+1}) &= 0, \\ \Rightarrow f(x_n) + \left(\frac{1}{2}\right)(x_{n+1} - x_n)[f'(x_n) + f'(x_{n+1})] &= 0 \\ \Rightarrow x_{n+1} &= x_n - \frac{2f(x_n)}{f'(x_n) + f'(x_{n+1})}. \end{aligned} \quad \dots(1.6)$$

This is an implicit scheme, which means derivative of the function at the $(n+1)^{th}$ iterative step is used to calculate the $(n+1)^{th}$ iterate. This difficulty is overcome by using Newton's method to compute the $(n+1)^{th}$ iterate on the right-hand side. Therefore, the resulting scheme is

$$x_{n+1} = x_n - \frac{2f(x_n)}{[f'(x_n) + f'(x_{n+1}^*)]}, \quad n=0,1,2,\dots \quad \dots(1.7)$$

$$\text{where } x_{n+1}^* = x_n - \frac{f(x_n)}{f'(x_n)}. \quad \dots(1.8)$$

1.3 Convergence of Method

Let α be a simple root of $f(x)$, i.e., $f(\alpha)=0$ and $f'(\alpha)\neq 0$. Let approximate value of the root is given by $x_n = \alpha + e_n$, where e_n is the error. Using Taylor expansion $f(x_n)$ can be written as:

$$\begin{aligned} f(x_n) &= f(\alpha + e_n) \\ &= f(\alpha) + f^{(1)}(\alpha)e_n + \frac{1}{2!}f^{(2)}(\alpha)e_n^2 + \frac{1}{3!}f^{(3)}(\alpha)e_n^3 + O(e_n^4) \\ &= f^{(1)}(\alpha) \left[e_n + \frac{1}{2!} \frac{f^{(2)}(\alpha)e_n^2}{f^{(1)}(\alpha)} + \frac{1}{3!} \frac{f^{(3)}(\alpha)e_n^3}{f^{(1)}(\alpha)} + O(e_n^4) \right] \\ &= f^{(1)}(\alpha) \left[e_n + C_2 e_n^2 + C_3 e_n^3 + O(e_n^4) \right], \end{aligned} \quad \dots(1.9)$$

where $C_j = (1/j!) f^{(j)}(\alpha) / f^{(1)}(\alpha)$. Similarly, using Taylor expansion,

$$\begin{aligned} f^{(1)}(x_n) &= f^{(1)}(\alpha + e_n) \\ &= f^{(1)}(\alpha) \left[1 + 2C_2 e_n + 3C_3 e_n^2 + O(e_n^3) \right] \end{aligned} \quad \dots(1.10)$$

Dividing equation (1.9) by equation (1.10) and after some simplifications, one gets,

$$\frac{f(x_n)}{f^{(1)}(x_n)} = e_n - C_2 e_n^2 + (2C_2^2 - 2C_3)e_n^3 + O(e_n^4), \quad \dots(1.11)$$

Substituting the results from equation (1.11) in equation (1.8), one gets,

$$x_{n+1}^* = \alpha + C_2 e_n^2 + (2C_3 - 2C_2^2)e_n^3 + O(e_n^4). \quad \dots(1.12)$$

This value of x_{n+1}^* is used for the Taylor's series expansion of $f^{(1)}(x_{n+1}^*)$ as,

$$\begin{aligned} f^{(1)}(x_{n+1}^*) &= f^{(1)}(\alpha) + [C_2 e_n^2 + (2C_3 - 2C_2^2)e_n^3 + O(e_n^4)] f^{(2)}(\alpha) + O(e_n^4) \\ &= f^{(1)}(\alpha) \left[1 + 2C_2^2 e_n^2 + 4C_2(C_3 - C_2^2)e_n^3 + O(e_n^4) \right] \end{aligned} \quad \dots(1.13)$$

Adding equation (1.10) and equation (1.13),

$$f^{(1)}(x_n) + f^{(1)}(x_{n+1}^*) = 2f^{(1)}(\alpha) \left[1 + C_2 e_n + \left(C_2^2 + \frac{3}{2}C_3 \right) e_n^2 + O(e_n^3) \right] \quad \dots(1.14)$$

using equation (1.9) and equation (1.14), one gets,

$$\begin{aligned} \frac{2f(x_n)}{[f^{(1)}(x_n) + f^{(1)}(x_{n+1}^*)]} &= \\ [e_n + C_2 e_n^2 + C_3 e_n^3 + O(e_n^4)] \times \left[1 + C_2 e_n + \left(C_2^2 + \frac{3}{2}C_3 \right) e_n^2 + O(e_n^3) \right]^{-1} & \end{aligned}$$

on simplifying it becomes, $\frac{2f(x_n)}{[f^{(1)}(x_n) + f^{(1)}(x_{n+1}^*)]} = e_n - \left(C_2^2 + \frac{1}{2}C_3 \right) e_n^3 + O(e_n^4).$

... (1.15)

Therefore from equation (1.7) and equation (1.15)

$$\begin{aligned}
 x_{n+1} &= x_n - \frac{2f(x_n)}{f^{(1)}(x_n) + f^{(1)}(x_{n+1}^*)}, \\
 e_{n+1} + \alpha &= e_n + \alpha - \left[e_n - \left(C_2^2 + \frac{1}{2} C_3 \right) e_n^3 + O(e_n^4) \right], \\
 e_{n+1} &= \left(C_2^2 + \frac{1}{2} C_3 \right) e_n^3 + O(e_n^4). \tag{1.16}
 \end{aligned}$$

This shows that the proposed method has third-order convergence.

1.4 Numerical Examples

The authors have demonstrated the use of their variant of Newton's method over the classical Newton's method for the examples given in Table 1.1. The roots were found correct to 15 decimal places.

1.5 Conclusions

From Table 1.1 it was found that the computational order of convergence is more than three in some cases in variant of Newton's method, which is higher than the classical Newton's method. The number of function evaluations was found to be less for variant of Newton's method as compared to classical Newton's method. Another important characteristic of this method is that it does not require second or higher derivatives of the function to carry out iterations.

Table 1.1 : Examples used for comparison of Variant's Newton method and classical Newton's method

S.No.	Functions	Root
1.	$x^3 - 4x^2 - 10 = 0$	1.36523001341448
1.	$\sin^2 x - x^2 + 1 = 0$	1.40449164821621
3.	$x^2 - e^x - 3x + 2 = 0$	0.25753028543977
4.	$\cos x - x = 0$	0.73908513321475
5.	$(x - 1)^3 - 1 = 0$	1.000000000000000
6.	$x^3 - 10 = 0$	1.15443469003367
7.	$xe^{x^2} - \sin^2 x + 3\cos x + 5 = 0$	-1. 20764782713013
8.	$x^2 \sin^2 x + e^{x^2 \cos x \sin x} - 28 = 0$	4.82458931731526
9.	$e^{x^2+7x-30} - 1 = 0$	3.000000000000000

REFERENCES

- [1] H. Brézis, Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert, North Holland Mathematics Studies, vol. 5, North-Holland, Amsterdam, London, 1973.
- [2] L. Collatz, Functional Analysis and Numerical Mathematics, Translated from the German by Hansjörg Oser, Academic Press, New York, 1966.
- [3] K. Deimling, Nonlinear Functional Analysis, Springer Verlag, Berlin, Heidelberg, New York, 1985.
- [4] V. Hutson and J. S. Pym, Applications of Functional Analysis and Operator Theory, Academic Press, New York, 1980.
- [5] L. V. Kantorovich, On Newton's method, Trudy Mat. Inst. Steklov. 28 (1949), 104– 144.
- [6] L. V. Kantorovich and G. Akilov, Functional Analysis. Transl. from the Russian, Pergamon Press, Oxford, 1982.

- [7] S. G. Krantz and H. R. Parks, The Implicit Function Theorem. History, Theory and Applications, Birkhäuser Verlag, Boston, Basel, Berlin, 2002.
- [8] J. W. Neuberger, A Nash-Moser theorem with near-minimal hypothesis, Int. J. Pure Appl. Math. 4 (2003), 269–280.
- [9] C. T. Rajagopal and T. V. Vedamurti Aiyar, A Hindu approximation to π , Scripta Math. 18 (1952), 25–30.
- [10] R. Rashed, The Development of Arabic Mathematics: Between Arithmetic and Algebra, Kluwer, Dordrecht, The Netherlands, 1994.
- [11] J. Schröder, Über das Newtonsche Verfahren, Arch. Rational Mech. Anal. 1 (1957), 154–180.
- [12] D. T. Whiteside (ed.), The Mathematical Papers of Isaac Newton, Volume II, 1667- 1670, Cambridge University Press, Cambridge, 1968.
- [13] D. T. Whiteside (ed.), The Mathematical Papers of Isaac Newton, Volume III, 1670-1673, Cambridge University Press, Cambridge, 1969.
- [14] T. J. Ypma, Historical development of the Newton-Raphson method, SIAM Review 37 (1995), 531–551.

