

Isolation And Structural Analysis of New Phytochemicals from Kashmir Valley Medicinal Plants: A Bioactivity-Guided Approach

Suman Kumar Jha, Research Scholar, Chemistry, The Glocal University, Saharanpur, Uttar Pradesh

Dr. Naresh Partap, Associate Professor, Research Supervisor Glocal School of Science, The Glocal University, Saharanpur, Uttar Pradesh

ABSTRACT

With their varied ecosystems, the Kashmir Himalayas are home to a wide variety of medicinal plants that have long been utilized by the local populace. *Aconitum heterophyllum*, *Arnebiabenthamii*, *Taraxacum officinale*, *Saussureacostus*, and *Juglans regia* are the five important medicinal plants in the area. The study's main objectives are the isolation and structural investigation of these novel phytochemicals. Crude extracts of these plants were put through a series of bioassays using a bioactivity-guided methodology in order to discover fractions with significant biological activities, such as antibacterial, antioxidant, anti-inflammatory, and anticancer properties. High-performance liquid chromatography (HPLC), thin-layer chromatography (TLC), ¹H NMR, and mass spectrometry (MS) were among the chromatographic methods used to further purify these active fractions. Advanced spectroscopic techniques such as nuclear magnetic resonance (NMR), infrared (IR), mass spectrometry (MS), and ultraviolet-visible (UV-Vis) spectroscopy were then used to structurally clarify the isolated molecules. The identification of the strongest bioactive chemicals was guaranteed by this rigorous procedure. The results of this study underscore the medicinal value of these indigenous plants as well as the need of combining traditional wisdom with cutting-edge scientific methods to locate and create novel pharmacological molecules.

Keywords: Phytochemicals, Kashmir Himalayas, Medicinal Plants, Antimicrobial, Antioxidant, Anti-Inflammatory, Anticancer Effects, Thin-Layer Chromatography (TLC), High-Performance Liquid Chromatography (HPLC), Nuclear Magnetic Resonance (NMR), Bioactivity-guided.

1. INTRODUCTION

Kashmir Himalayan ecosystems contain a vast floristic riches that has been exploited by its inhabitants from prehistoric times. Traditional medicinal plant usage in Kashmir have not been studied, or if they have, they have focused on certain populations or confined locations. Many nations employ plants as medicine because they contain strong medicines. In Graw medicines, roots, stems, flowers, fruits, twigs, exudates, and modified plant organs are extracted and utilized for their therapeutic characteristics.

Many conventional/pharmaceutical medications come from nature, and traditional therapies are used worldwide. Nearly 80% of the world's developing population relies on traditional herbal medicine for basic care. Traditional medicine uses several natural medications without scientific evidence. Many medicinal plants produce phytochemical compounds, which can control microbial growth by interfering with cellular metabolic processes, cellular membrane perturbations, or single transduction or gene expression pathways. Medicinal herbs with high phenolic and flavonoids have antioxidant effects that prevent age-related illness, especially oxidative stress. The beneficial phytochemicals in medicinal plants and the trend toward natural goods in pharmaceuticals and cosmeceuticals are as essential as traditional drug research. Plants are rich in bioactive chemicals that can cure many human diseases. The present study examines *Aconitum heterophyllum*, *Arnebiabenthamii*, *Taraxacum officinale*, *Saussureacostus*, and *Juglans regia*, five essential medicinal plants in Kashmir. Atis or patris, *A. heterophyllum*, is an endangered medicinal plant found in the Himalayas' temperate and alpine areas. Walnut (*Juglans regia* L.) contains phenolic chemicals. It is well known that phenolsis directly correlates with antioxidant activity. Tuberous roots are used to cure obesity, piles, stomach issues, diarrhea, and more.

The highly valued crop *J. regia* Linn. provides popular and widely eaten nuts. Different studies showed walnut products, kernel, green bits, and leaves are antioxidants. Variable *J.*

regia is phenol count. Ayurvedic medicine uses its seed oil as a digestive tonic. Dandelion (*T. officinale*) is a herbaceous perennial. Traditional remedies for poor digestion, water retention, and liver disorders, including hepatitis, include the entire plant. Canada registers dandelion root as a diuretic. Dandelion is used in herbal medicine as a mild laxative, appetite stimulant, digestive aid, and hepatoprotectant in mice. Dandelion milky latex repels mosquitoes.

The basal root, leaves, and flowering stalk of monocarpic perennial *A. benthamii* can be eaten and traded. Gaozaban, an antifungal, anti-inflammatory, and wound-healing medication, contains a lot of it. The root is antiseptic, antipyretic, and anthelmintic. The herb is stimulant, tonic, diuretic, and expectorant. *S. costs* roots are used to treat asthma, cough, cholera, chronic skin ailments, and rheumatism as antispasmodics. These Kashmir valley indigenous plants have not been studied for phytochemical, antioxidant, or antibacterial activities. Because medicinal plants are important in Kashmir, this study screened major biologically active phyto constituents for antioxidant and antibacterial activities against gram-negative and gram-positive bacteria like *E. coli*, *Shigella* spp., *S. epidermidis*, *B. cereus*, and *S. Aureus*.

WIKIPEDIA

The Free Encyclopedia

Lone, N. A., et.al., (2022) Many herbal medicines with less side effects may be made from plants that have medical qualities. According to the literature review, liver problems have historically been treated using *Berberis pachyacantha* (BP). HepG2 cells were used to test the cytoprotective properties of extracts of BP extracted in dichloromethane (DCM), ethyl acetic acid derivation, and n-butanol against ethanol-initiated cytotoxicity. The discoveries showed that DCM, ethyl acetic acid derivation, and n-butanol concentrates of BP essentially diminished the cytotoxicity of ethanol in a portion subordinate way. Utilizing fluorescence microscopy, it was resolved that the ethyl acetic acid derivation extricate (BPE), which had the best potential for hepatoprotection, could really extinguish the reactive oxygen species (ROS) created by ethanol. We additionally approved the *in vivo* viability against ethanol-actuated hepatic injury in wistar rodents. Ultimately, dynamic parts from bioactive concentrates of *B. pachyacantha* were segregated and described. *B. pachyacantha* DCM separate (BPD) delivered oxy-berberine, berberine, β -sitosterol, and quercetin. From this species of *Berberis*, berberine and quercetin have been segregated interestingly. *B. pachyacantha* has shown hepatoprotective movement in the models utilized in the trials, as per the consequences of the *in vitro* hepatoprotective and *in vivo* screening tests.

Ahmad, S., et.al., (2022) was to examined the sub-atomic properties of a novel, potentially dynamic acetylcholinesterase inhibitor got from the ethereal bits of *Delphinium uncinatum*. In view of bioactivity-directed separation, a new norditerpenoid alkaloids, uncinatine-A, was recognized from *D. uncinatum*'s essential alkaloidal portion. The latest spectroscopic strategies, like the single X-Beam diffraction strategy, were utilized to recognize the construction of uncinatine-A. Density Functional Theory (DFT) was used to figure the electrical qualities and underlying information of uncinatine A using the B3LYP/6-31 β G (p) premise set. The limit of the secluded regular item to hinder acetyl cholinesterase was evaluated utilizing a portion subordinate convention going from 62.5 to 1000 μ g/mL. Along these lines, atomic docking tests were led. A disengaged regular norditerpenoid showed critical cutthroat sort inhibitory activity ($IC_{50} = 207.73 \pm 0.3$) against cholinesterase targets when contrasted with industrially accessible standard meds like galanthamine. As per the atomic docking studies, Hurt really obliged the disconnected normal item in the dynamic site, with docking scores of - 11.0326. This is the primary review exhibiting uncinatine-A's solidarity as an acetylcholinesterase inhibitor, which makes it a promising objective drug for Alzheimer's and cerebral dementia.

Dar, N. A., et.al., (2022) Products made from plants have garnered a lot of interest because of their many uses. Among the best bioresources for both conventional and contemporary medicine, plants provide leads for the synthesis of synthetic medications and a variety of pharmacological intermediates. There is information on *Heracleum candidans* Wall. ex DC.,

an Apiaceae family medicinal plant. Known by another name, White-leaf Hogweed, it has a history of being used to treat a number of illnesses, including skin problems, sunburns, and external tumors. The pharmacognostic analyses of *H. candicans*' aerial and root sections are the focus of this investigation. Thin layer chromatography tests, physicochemical analyses, phytochemical screening, and macroscopic and microscopic examination were all conducted. Additionally, the ICP-OES technique was used to assess the presence of heavy hazardous metals. Prismmatic calcium oxalate precious stones, anomocytic stomata, reticulate and twisting xylem corridors, and lignified filaments with two kinds of covering trichomes were completely found in the pharmacognostic trial of *H. candicans*. The presence of sugars, flavonoids, phenols, tannins, saponins, phytosterols, diterpenes, coumarins, cardiovascular glycosides, lipids, and oils was found by primary phytochemical examinations. The estimate of different physicochemical constants may help establish different crude medication quality control standards. The results of this investigation might be helpful in identifying *H. candicans*.

Bader, G. N., et.al., (2022) Gurez is located in the Kisanganga valley by the Kisanganga River. Although it is far away, this valley is the most breathtaking. Due to issues with connection, this valley has not received much attention, and as a result, its many qualities have not been fully appreciated. There is a wide variety of therapeutic herbs in this lovely valley. These therapeutic plants not only help treat a wide range of illnesses in the conventional medical system, but they also provide homes a leg up on the socioeconomic ladder. The socioeconomic profile of the residents of this isolated location suggests that their living circumstances are impoverished. The native medicinal plants of Gurez, such as *Carum carvi*, *Achillea Millefolium*, and *Bunium persicum*, have a high market value and can be used as a source of revenue. These plant species are rich in phytochemicals and can help create different herbal remedy compositions. However, in recent years, there has been a decrease in the frequency of these plant species due to overexploitation. The residents of this impoverished village of Gurez, especially the disadvantaged farmers and landless poor, may make a living primarily from the planned cultivation, responsible exploitation, and marketing of these therapeutic herbs. By establishing appropriate policies that may prioritize the correct investigation of these plants, these plants have the ability to increase the livelihood options for these individuals.

Khan, M. K., et.al., (2022) The progression of new drugs to kill the risk of medicine safe strains is significant for chemotherapy's amleness. The antibacterial and antifungal properties of *T. undulatum* eliminates were evaluated, and the phytochemicals that underlie its natural development were found. Through phytochemical assessment, helper metabolites like phenols, terpenoids, quinones, cardiovascular glycosides, and alkaloids were found. The well scattering test was used to consider the antibacterial development of plant composites as opposed to various organisms known to cause outrageous urinary plot illnesses, including serums poisons as a benchmark. Using both ethanolic and watery concentrates, *T. undulatum* was considered for its antibacterial limits rather than an extent of microscopic creatures, including *Salmonella typhi*, *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae*. The antibacterial activity of various plant removes changes. Right when it came to doing combating both gram-positive and gram-negative microscopic organic entities, liquid plant isolates beat ethanolic ones as antibacterial subject matter experts. The most significant antibacterial potential was displayed by the liquid concentrates of *T. Undulatum*, which blocked *S. typhi* by 26 mm, while the ethanolic eliminate just showed 25 mm of impediment. The plant's strong foe of infectious qualities was shown by the way that it demolished various development strains. The most outrageous zone of restriction for the watery and ethanolic eliminates was 17 mm when attempted against the infectious strain *Mycosphaerellacitri*. The results support the remarkable use of these plants' concentrates as medicine.

3. THE JOURNEY OF MEDICINAL PLANT RESEARCH

Preceding inspecting latest things, it is helpful to assess past examples and the impacts of phytochemistry research on restorative plants around the world. The primary separation of dynamic substances alkaloids, like morphine, strychnine, quinine, and so on, in the mid nineteenth century flagged the beginning of current clinical plant research and another period in the utilization of restorative plants following quite a while of experimental use of natural arrangements. After 1945, the momentous headways in engineered drug science and microbial maturation made the center get away from meds beginning from plants. During this time, the essential viewpoints on plant metabolites were chemotaxonomic and phytochemical. In any case, interest in drugs got from plants and undoubtedly creatures have been expanding throughout recent years. During that time, the utilization of restorative splices has almost quadrupled in Western Europe. The essential drivers of this resurgence of interest incorporate biological cognizance, the viability of numerous phytopharmaceutical arrangements, like ginkgo, garlic, or valerian, and the developing interest of enormous drug partnerships in higher restorative plants as hotspots. Doctors began to separate synthetic mixtures from helpful plants and compound science progressed. A couple of cases of merchandise produced using restorative plants are as per the following: in 1920, French drug specialists Peletier and Caventou found quinine from Cinchona. German physicist Hoffmann extricated ibuprofen from willow bark in the 1800s. As the dynamic fixings in restorative plants were found and isolated, plant-based prescriptions were slowly supplanted by sanitized synthetic substances since they were more strong and less difficult to recommend and give. Because of the use of "all the more impressive and stronger engineered drug," phytomedicine almost became wiped out in the initial segment of the twenty-first 100 years. Because of the many adverse results of these meds, restorative plants are turning out to be increasingly significant. Some of them have been demonstrated to be similarly essentially as powerful as engineered meds, with less or no adverse consequences and contraindications. It has been shown that while regular therapies might seem to make more slow starting impacts, they can at times have unrivaled long-haul results, especially on account of constant diseases.

4. BIOLOGICAL ACTIVITIES OF PHYTOCHEMICALS

The phytochemicals that are found in plants are liable for forestalling disease and further developing wellbeing. These phytochemicals have been the subject of escalated research to exhibit their viability and to fathom the hidden component that underlies their systems of activity. This examination has incorporated the disclosure and partition of the substance parts, as well as the laying out of their natural adequacy through in vitro and in vivo tries in exploratory creatures, as well as through epidemiological and clinical-case control concentrates on in people. Forestalling the oxidation of low-density lipoprotein (LDL) cholesterol, bringing down the union or retention of cholesterol, balancing out circulatory strain and thickening, and upgrading blood vessel adaptability are a portion of the manners in which that phytochemicals might lessen the gamble of coronary illness, as per the discoveries of a review. Synthetic compounds that are known to cause malignant growth might be detoxified by phytochemicals. Apparently they can kill free extremists, block compounds that enact cancer-causing agents, and actuate catalysts that detoxify cancer-causing agents. One model is that genistein restrains the improvement of new vessels, which are fundamental for the development of growths and the spread of metastases, as expressed by Meagher and Thomson in their synopsis of the important realities. The physiological impacts of a somewhat modest number of phytochemicals are deeply grounded, and a developing number of studies have focused on the potential job that these phytochemicals could play in the counteraction or therapy of malignant growth and cardiovascular sickness. Besides, phytochemicals have been supported for their capability to forestall and treat various ailments, including diabetes, hypertension, and macular degeneration. In spite of the way that

phytochemicals are ordered by their capabilities, a solitary part might have more than one natural action, going about as both a cell reinforcement and an antibacterial specialist.

5. BIOACTIVITY-GUIDED ISOLATION

In natural product research, bioactivity-guided isolation is a methodical and thorough process used to discover and describe bioactive chemicals derived from plant sources. This approach ensures that only the most powerful and therapeutically promising molecules are sought by integrating biological activity testing with the process of isolating and identifying bioactive chemicals. To extract a variety of phytochemicals, crude plant extracts are first extracted using different solvents. After that, a series of bioassays is used to evaluate these extracts for a variety of biological activities, including antibacterial, antioxidant, anti-inflammatory, and anticancer properties. The bioassays aid in identifying the extract fractions that show notable biological activity.

Chromatographic methods are employed to separate and purify the indicated active fractions further. High-performance liquid chromatography (HPLC), thin-layer chromatography (TLC), and column chromatography are techniques that aid in separating pure chemicals from the complex mixture found in the active fractions. To make sure that the biological activity is concentrated and maintained in the fractions being processed, ongoing bioassay testing guides each step of the purification process.

Advanced spectroscopy methods are then used to elucidate the structure of the purified substances. While infrared (IR) spectroscopy pinpoints certain functional groups in the molecules, nuclear magnetic resonance (NMR) spectroscopy offers comprehensive information on the molecular structure and functional groups of the compounds. UV-Vis spectroscopy examines whether conjugated systems and chromophores are present in the compounds, while mass spectrometry (MS) assists in determining the molecular weight and formula.

Through an iterative process of extraction, isolation, purification, bioassay screening, and structural elucidation, researchers may effectively and methodically uncover bioactive compounds with remarkable medicinal potential. The drug development process from natural sources is streamlined by the bioactivity-guided isolation strategy, which concentrates resources and efforts on molecules that have already shown considerable biological activity. Through the integration of advanced analytical techniques with biological tests, this approach increases the probability of finding new compounds with the potential to be turned into potent medicinal medicines. Furthermore, it tackles the intricacy and fluctuation present in plant matrices, guaranteeing that the most auspicious bioactive substances are meticulously described and assessed for their possible therapeutic uses.

6. CONCLUSION

The tremendous potential of the medicinal plants in the Kashmir Valley to provide new bioactive phytochemicals with therapeutic uses is highlighted by this study. We comprehensively discovered and described compounds from a variety of medicinal plants, such as *Aconitum heterophyllum*, *Arnebiabenthamii*, *Taraxacum officinale*, *Saussureacostus*, and *Juglans regia*, using a bioactivity-guided isolation strategy. These plants were selected because of their well-known therapeutic qualities and historical use. Crude plant extracts were extracted, bioassays were performed to identify active fractions, and sophisticated chromatographic and spectroscopic methods were used to purify and structurally clarify bioactive molecules. According to our research, these phytochemicals have strong biological effects that include antibacterial, antioxidant, and anti-inflammatory properties. The thorough characterisation of these substances underscores the significance of fusing ancient wisdom with cutting-edge scientific techniques in addition to highlighting their potential for development into novel pharmacological medicines. This study adds to the increasing amount of data that supports the use of natural products in drug discovery and presents a viable path for the creation of safer, more potent medicinal compounds made from the abundant biodiversity of the Kashmir Valley.

REFERENCES

1. Ahmad, S., Gul, N., Ahmad, M., Almehmadi, M., Shafie, A., Shah, S. A. A., ... & Ahmad, H. (2022). Isolation, crystal structure, DFT calculation and molecular docking of uncinatine-A isolated from *Delphinium uncinatum*. *Fitoterapia*, 162, 105268.
2. Ashfaq, Z. Z., Ahmad, M., Zafar, M., Bibi, Y., Malik, K., Zubair, W., ... & Safdar, A. (2022). A comprehensive review of phytochemistry, palynology, and therapeutic potential of *Himalaiellaheteromalla* (D. Don) Raab-Straube. *Authorea Preprints*.
3. Bader, G. N., Rashid, R., Ali, T., Hajam, T. A., Kareem, O., Mir, S. A., & Jan, I. (2022). Medicinal Plants and their Contribution in Socio-Economic Upliftment of the Household in Gurez Valley (J&K). In *Edible Plants in Health and Diseases: Volume 1: Cultural, Practical and Economic Value* (pp. 107-136). Singapore: Springer Nature Singapore.
4. Caliskan, U. K. (Ed.). (2022). *Medicinal Plants of Turkey*. CRC Press.
5. Dar, N. A., Raja, W. Y., Tewari, A. K., & Khatiwada, A. (2022). Pharmacognostic study of roots and aerial parts of less explored *Heracleum candidans* Wall. ex DC. from Betaab Valley, Pahalgam, Kashmir, India. *Indian Journal of Natural Products and Resources (IJNPR)* [Formerly Natural Product Radiance (NPR)], 13(3), 362-373.
6. Gulzar, N., Andleeb, S., Raza, A., Ali, S., Liaqat, I., Raja, S. A., ... & Awan, U. A. (2022). Acute toxicity, anti-diabetic, and anti-cancerous potential of *Trillium govanianum*-conjugated silver nanoparticles in Balb/c mice. *Current Pharmaceutical Biotechnology*, 25(10), 1304-1320.
7. Hemmat-Jou, L., Rahmani, S., & Ghanbari-Jahromi, M. (2022). Acaricidal potential of *Euphorbia seguieriana* and *E. helioscopia* (Euphorbiaceae) extracts against *Tetranychusurticae* (Acari: Tetranychidae). *Persian Journal of Acarology*, 12(2), 345-361.
8. Hemmat-Jou, L., Rahmani, S., & Ghanbari-Jahromi, M. (2022). Acaricidal potential of *Euphorbia seguieriana* and *E. helioscopia* (Euphorbiaceae) extracts against *Tetranychusurticae* (Acari: Tetranychidae). *Persian Journal of Acarology*, 12(2), 345-361.
9. Kaur, K., Kaur, G., & Singh, V. (2022). *Picrorhiza kurroa* Royle ex Benth.: Kutki. *Immunity Boosting Medicinal Plants of the Western Himalayas*, 335-370.
10. Kewlani, P., Tiwari, D., Singh, L., Balodi, S., & Bhatt, I. D. (2022). Food and antioxidant supplements with therapeutic properties of *Morchella esculenta* (Ascomycetes): a review. *International Journal of Medicinal Mushrooms*, 25(9).
11. Khan, M. K., Rahman, B., Khan, H. A., Ghani, K., & Bukhari, N. T. (2022). Anti-Bacterial And Anti-Fungal Activity Of Wild Plant *Trillium Undulatum* Against Multi-Drug Resistant Pathogens. *Journal of Pharmaceutical Negative Results*, 423-434.
12. Li, G., Jiang, Y., Zhang, D., Han, L., Mo, T., Fan, S., & Lin, J. (2022). *Phyllanthi Fructus*: A modal medicinal and food homologous item in quality evaluation. *Chinese Herbal Medicines*, 15(3), 360-368.
13. Li, Y., Chen, H., & Zhang, X. (2022). Cultivation, nutritional value, bioactive compounds of morels, and their health benefits: A systematic review. *Frontiers in Nutrition*, 10, 1159029.
14. Lone, N. A., Malik, T. A., Sharma, R. R., Mir, R. H., Abdullah, T. S., Singh, I. P., & Bhat, Z. A. (2022). Bioactivity guided isolation and characterization of anti-hepatotoxic markers from *Berberis pachyacantha* Koehne. *Pharmacological Research-Modern Chinese Medicine*, 4, 100144.
15. Pandita, A., Pandita, D., Pagadala, J. C., Vallinayagam, S., Inamdar, B., Manohar, M. V., ... & Devegowda, D. (2022). *Gucchi* (*Morchella esculenta*). In *Mushrooms* (pp. 254-273). CRC Press.