



Analyzing Forest Community Distribution, Structure, and Regeneration in the Northwestern Himalayan State of Himachal Pradesh's Shivalik Hills

Mohd. Umar, Research Scholar, Environmental Science, Glocal School of Science, The Glocal university U.P.
Dr. Nirmal Sharma (Associate Professor), Glocal School of Science, The Glocal university U.P.

Abstract

The Shivalik Hills in the Western Himalayas are a natural sanctuary with a rich biodiversity and cultural legacy, including medicinal plants crucial to local healthcare systems. However, urbanization, deforestation, and climate change threaten these plants. A comprehensive assessment of the flora is being conducted to preserve its herbal resources. The study uses fieldwork and statistical research to create a database of plant traits, historic use, and distribution patterns. The study also highlights the importance of medicinal plants in the Indian traditional medicine system Ayurveda, which promotes holistic health and wellbeing. The study also analyzes medicinal plant pharmacological effectiveness to understand their therapeutic potential. The findings underscore the need for conservation efforts to preserve the ecological integrity and medicinal legacy of the Shivalik Hills.

Keywords: Shivalik Hills, Western Himalayas, Biodiversity, Cultural Legacy, Medicinal Plants

1. INTRODUCTION

The Shivalik Hills, situated in the Western Himalayas, have a distinct geographical location, a varied climate, and an abundance of natural diversity and cultural heritage. Numerous medicinal plants that have been utilized for millennia by indigenous tribes are found in these hills and are essential to their traditional healthcare systems. But as the environment's delicate equilibrium is threatened by unchecked deforestation, rising urbanization, and climate change, the hills confront ever-greater problems. Because of human activities increasing their presence in natural regions, many plant species, including medicinal plants, are in danger of becoming extinct. By conducting a thorough investigation to identify and catalog the various medicinal plants present in the area, as well as by assembling a comprehensive database based on botanical characteristics, traditional uses, and distribution patterns through fieldwork and statistical data collection, this thesis seeks to document and conserve the herbal abundance of the Shivalik Hills.

1.1. History Of Medicinal Plants

Life first appeared on Earth around 4.5 billion years ago. Early human civilizations relied heavily on plants; in Iraq, herbal medicine dates back 60,000 years, while in China, it dates back 8,000 years. Herbs have been used medicinally by the Neanderthal people since 5000 BCE. Indian texts with thorough descriptions of medical disorders include the Yajurveda, Atharva Veda, and Rigveda. Books like Sushruti Samhita, Dhanwantri Nighantu, and Charak Samhita highlight the use of plants and polyherbal remedies to treat a variety of illnesses. Consequently, the rich traditions and scientific legacies of past civilizations have led to the development of herbal treatments.

Ayurveda, Siddha, and Unani are the three established traditional medicinal systems in India. The use of plants for religious purposes, medicinal purposes, food production, fuel production, and agricultural equipment is mentioned in the Rigveda, Atharvaveda, Upanishads, Mahabharata, and Puranas. All three Samhitas (Sushruta, Charaka, and Ashtanga Hridaya) include more than 1200 plant-based medications, their effects, and how to use them in therapeutic contexts. Plant identities are unknown. Natural resource protection is the responsibility of indigenous peoples across the globe. Because of modern living and technology breakthroughs, traditional knowledge is quickly disappearing. These groups include approximately 250 million individuals, while the Indian subcontinent is home to 53 million tribal people from over 550 tribes. Tribes are discrete ethnic groupings that possess unique customs, traditions, food preferences, religious practices, and medical expertise. These tribes comprise 7.7% of India's total population.





1.2. Ayurvedic Medicine System

India has a long history of using traditional medicine, mostly based on Ayurveda, which is said to be the oldest medical system in the world. The science of life, or ayurveda, is regarded as one of the earliest forms of holistic medicine in India. It highlights how crucial it is for a person's physical, mental, and spiritual wellbeing to remain in balance. Many medicinal systems, such as Siddha, traditional Chinese medicine, and Unani medicine, have inspired Ayurveda. Following the conquest of India by Western colonial forces in the 18th and 19th centuries, traditional Indian culture was outlawed or suppressed. However, there has been a global revival of interest in Ayurvedic medicine due to the fact that more and more people are turning to it as a holistic, natural approach to health and wellbeing, and scientific evidence is beginning to support its efficacy.

2. LITERATURE REVIEW

Shah (2019) examines the background and applicability of Ayurveda, an ancient Indian medical system, worldwide. It highlights Ayurveda's comprehensive approach to health and healing and traces its origins back over five millennia to ancient India. The body, mind, and spirit are interrelated, and this connection serves as the basis for the system's diagnostic and therapeutic approaches. The study also emphasizes how ancient knowledge, embodied in books like as the Sushruta Samhita and Charaka Samhita, served as authoritative sources of wisdom in the development of Ayurvedic procedures. The assessment also emphasizes how Ayurveda has spread around the world and been embraced by many cultural settings, underscoring its growing appeal and acceptability. The review provides a detailed overview of the development, tenets, and current significance of Ayurveda by synthesizing available research and academic literature.

Pandey et al., (2013) The Indian Traditional Ayurvedic System of Medicine emphasizes the value of preserving harmony and balance within the body via a variety of treatment approaches. It takes a holistic approach to health and wellbeing. The review delves into the function of nutritional supplements in the context of Ayurveda, emphasizing the usage of minerals, herbs, and other natural substances to enhance and maintain overall health. The selection and administration of dietary supplements are informed by Ayurvedic principles, which take into account an individual's imbalances (vikriti) and constitution (prakriti). The study also looks at scientific evidence, based on both modern research results and traditional knowledge, that supports the efficacy and safety of Ayurvedic nutritional supplements. The combination of nutritional supplements and Ayurvedic therapy has the potential to improve health and well-being.

Adhikari and Paul (2018) explores the evolution of Indian traditional medicine, emphasizing the importance of its culture and heritage. It looks at how these traditions have changed throughout centuries, emphasizing the contributions of scholars, practitioners, and historic texts. The evaluation also touches on the passing down of medical practices and information from one generation to the next, emphasizing the resilience and continuity of India's traditional medical systems. The writers also discuss the role of Indian traditional medicine in modern healthcare, looking at how it might be incorporated with conventional methods and how it could improve health throughout the world. With its thorough examination of the rich history and current expansion of Indian Traditional Medicine, the book is an invaluable tool for scholars and medical professionals.

Raju et al., (2022) examines the pharmacological and therapeutic efficacy of Achyranthes aspera Linn, a plant that has been used historically in conventional medical systems. Anti-inflammatory, analgesic, antibacterial, antioxidant, and anti-cancer effects of the plant have been reported. In order to comprehend the active components in charge of the plant's bioactivity, the review synthesizes the results of earlier studies. Achyranthes aspera Linn has been used to treat a number of illnesses, such as inflammatory problems, cancer, and infectious infections. The evaluation also emphasizes dosage problems and the need for



uniform preparations. The paper gives a thorough summary of the plant's medicinal potential and suggests future avenues for drug development and medical research.

3. RESEARCH METHODOLOGY

3.1. Sample Size

The number of communities or community types that are mentioned in the data is referred to as the sample size. Four community kinds are present in this instance: mixed Acacia catechu-Cassia fistula, mixed Acacia catechu-Chamaerops humilis, and mixed Acacia catechu-Mallotusphilippensis.

3.2. Identification of species

The Shannon's Diversity Index (H'), which is a representation of the species diversity within each community type, is provided by the data. For every kind of community, the densities of different vegetation components, including trees, shrubs, seeds, and herbs, are also given. Nevertheless, the data does not include the names of the particular species. Rather, within each community type, the data conceives the densities and variety of plant components.

3.3. Research Design

The Free Encyclopedia

With an emphasis on describing and contrasting various community types according to their ecological and vegetational properties, the study strategy seems to be observational and descriptive. The goal of the research is to comprehend the vegetation dynamics and biodiversity found in Himachal Pradesh's Shivalik Hills.

3.4. Data Collection Techniques

- **Primary Data Collection**

Primary data gathering procedures are probably used for the data shown in "Table 1: Community Data Summary" and "Community Type Data (Table 2)". This contains measurements, ecological evaluations, and in-person field surveys carried out in Himachal Pradesh's Shivalik Hills. Through direct observations, field surveys, and ecological assessments carried out in the study area, researchers may have gathered data on total species, tree density, total basal area, species richness, altitudinal range, latitude range, longitude range, habitat characteristics, slope range, aspect, and major associates.

- **Secondary Data Collection**

The data shown in "Community Vegetarian Density (Table 3)" and "Figure 1: Graphical Representation on the Community Data" may have been gathered using a mix of primary and secondary sources. Geographical coordinates and habitat characteristics may have been obtained directly from field surveys and observations, while other data, such as species diversity (as determined by Shannon's Diversity Index, H') and vegetation densities, may have been gathered from ecological databases, previously published works, or studies carried out in comparable ecosystems or geographic areas. In addition to providing more context and insights into the vegetation dynamics and biodiversity patterns within the research region, this secondary data complements the original data gathered for the study.

4. DATA ANALYSIS



ADVANCED SCIENCE INDEX

The information supplied sheds light on the ecological traits of five different communities that are located within the research region. The total species richness, total tree density, and total basal area of each community are used to draw boundaries, giving an overview of the structure and composition of the vegetation in these ecosystems. With 150 species, Community 3 has the most overall species count, making it stand out as a rich biodiversity hotspot. With just 25 species, Community 2 on the other hand, has the least variety of species. With 155 trees per unit area, Community 3 tops the list of areas with the highest tree density, indicating a thick forest canopy. Community 5, on the other hand, has the lowest tree density, suggesting a minimal amount of plant cover in its habitat. The vegetation structure is further clarified by total basal area measurements. Community 5 has the largest basal area while having a lower tree density, which may indicate that bigger, more mature trees are more common there. On the other hand, Community 3 has a somewhat smaller total basal area in

spite of its high tree density, suggesting that a combination of young and old trees contribute to the canopy. All things considered, these data provide insightful information on the variety, density, and organization of plant communities in the research region, which helps to clarify the ecological dynamics and preservation requirements of these ecosystems.

Table 1: Summary of Community Data




Figure 1: Graphical Representation on the Community Data

Table 2: Communities in the Shivalik Hills of Himachal Pradesh: kinds, distribution, and important associations.

Community type	S R	Altitudinal Range	Latitude Range	Longitude Range	Habitat(s)	Slope Range	Aspect(s)	Major Associates
Acacia catechu	15	350-599	30.35-32.00	76.00-77.52	Shady Moist, Rocky, Dry	5-30	E, N, W, NE, NW, SE, SW	Mallotusphilip pensis, Dalbergia sissoo, Cassia fistula
Acacia catechu- Cassia fistula mixed	3	710-765	25.78-25.69	71.41-71.20	Shady Moist, Dry	10-35	SE	Mallotusphilip pensis, Dalbergia sissoo, Albizia lebbeck
Acacia catechu- Chamaeropshumilis mixed	2	485-450	31.50-31.89	56.26-76.60	Riverine, Dry	5-25	NE	Dalbergia sissoo, Cassia fistula, Albizia lebbeck
Acacia catechu- Mallotusphilip pensis mixed	3	615-750	31.55-32.05	76.10-76.65	Shady Moist, Dry	10-30	W, SE, NSE	Dalbergia sissoo, Cassia fistula

Four unique community types are identified by the data presented, and each is distinguished by a range of ecological features. The first community type, Acacia catechu, is found in altitudinal ranges of 350–599 meters and has a high species richness (SR) of 15. Its range of



habitat includes dry, rocky, shady, and damp areas with slopes between 5 and 30 degrees and east, north, west, northeast, northwest, southeast, and southwest faces. *Cassia fistula*, *Dalbergia sissoo*, and *Mallotusphilippensis* are important allies in this community. In contrast, the *Acacia catechu*-*Cassia fistula* mixed community inhabits altitudinal regions of 710-765 meters and has a lower SR of 3. Its preferred habitats are shaded, damp, and dry, with slopes between 10 and 35 degrees, mostly facing southeast. *Albizia lebbeck*, *Dalbergia sissoo*, and *Mallotusphilippensis* are important allies in this community. Similarly, the mixed community of *Acacia catechu*-*Chamaeropshumilis* has altitudinal ranges of 485-450 meters and an SR of 2. Primarily found on northeast-facing slopes in riverine and arid settings, *Dalbergia sissoo*, *Cassia fistula*, and *Albizia lebbeck* are its main partners. Finally, the mixed community of *Acacia catechu*-*Mallotusphilippensis* has an SR of 3 and is found in altitudinal ranges between 615 and 750 meters. Its range of habitat includes dry, damp, and shaded areas with slopes between 10 and 30 degrees with faces that include the west, southeast, and north-south. Two important members of this group are *Cassia fistula* and *Dalbergia sissoo*. All things considered, these community types reflect a variety of biological niches, each suited to certain environmental circumstances and distinguished by distinct species compositions and habitat preferences.


Table 3: The recognized groups' structural structure in the Shivalik highlands of Himachal Pradesh

Community Type	Trees				Shrubs	Herbs
	Density	TBA (m ² /ha ⁻¹)	Saplings Density (Ind/ha ⁻¹)	Seedlings Density (Ind/ha ⁻¹)	Density (Ind/ha ⁻¹)	Density (Ind/ha ⁻¹)
Acacia catechu	420.65	15.08	450.50	166.50	1450.10	25.65
Acacia catechu-Cassia fistula mixed	600.00	19.72	260.00	640.00	915.00	25.10
Acacia catechu-Chamaeropshumilis mixed	430.00	5.47	565.00	810.00	1275.00	45.85
Acacia catechu-Mallotusphilippensis mixed	250.00	44.35	254.45	501.45	1445.10	25.50

With regard to tree density, shrub density, herb density, total basal area (TBA) per hectare, saplings density (individuals per hectare), and seedlings density (individuals per hectare), the data presented outline the features of four different kinds of communities. There are 420.65 trees per hectare in the *Acacia catechu* community, along with 15.08 shrubs and 450.50 herbs. The whole basal area is 166.50 square meters per hectare, and the density of seedlings and saplings is especially high at 25.65 and 1450.10 individuals per hectare, respectively. Comparably, the *Acacia catechu*-*Cassia fistula* mixed community has 600.00 trees per hectare of thick tree cover, with an additional 19.72 shrubs and 260.00 herbs. With sapling density of 915.00 individuals per hectare and seedling density of 25.10 individuals per hectare, TBA achieves 640.00 square meters per hectare. The mixed community of *Acacia catechu* and *Chamaeropshumilis* has a tree density of 430.00 individuals per hectare, 5.47 shrubs, and 565.00 herbs. TBA is 810.00 square meters per hectare, and its sapling and seedling densities are also quite high at 1275.00 and 45.85 individuals per hectare, respectively. Last but not least, the mixed community of *Acacia catechu* and *Mallotusphilippensis* has fewer trees—250.00 individuals per hectare—but more shrubs—44.35 individuals and 254.45 herbs—to

make up for it. With seedlings at 25.50 individuals per hectare and saplings at 1445.10 individuals per hectare, TBA reaches 501.45 square meters per hectare. Overall, representing differences in vegetation density and structure, these data provide insights into the ecological processes and composition of each community type.

Table 4: Concentration of Dominance (Cd) by community in the Shivalik highlands of Himachal Pradesh.



The data supplied shows the densities of different vegetation components in four distinct community types, as well as the species diversity (as determined by Shannon's Diversity Index, H'). With tree density at 1.22 individuals, saplings at 0.15 individuals, seedlings at 1.51 individuals, shrubs at 2.77 individuals, and herbs unidentified, the species diversity (H') in the Acacia catechu community is 0.45. Higher species diversity (H') of 1.35 is shown by the Acacia catechu-Cassia fistula mixed community, where tree density is 2.15 individuals, saplings are 0.45 individuals, seedlings are 1.77 individuals, shrubs are 2.51 individuals, and herbs are not defined.

Species diversity (H') for the mixed community of Acacia catechu-Chamaerops humilis is 1.65; tree density is 2.02 individuals, saplings are 0.22 individuals, seedlings are 2.22 people, shrubs are 3.22 individuals, and herbs are not given. The species diversity (H') in the mixed community of Acacia catechu and Mallotusphilippensis is 1.45. Tree density is 1.77 individuals, saplings are 0.45 individuals, seedlings are 2.45 individuals, shrubs are 2.65 individuals, and herbs are not specified. All things considered, these data illustrate differences in vegetation density and structure across various ecosystems and provide information on the species variety and makeup of each community type.

5. CONCLUSION

The Shivalik Hills' ecological landscape and medicinal flora have been studied extensively, providing valuable insights into the region's biodiversity and cultural heritage. The study reveals distinct ecological niches with varying species richness, vegetation density, and structural patterns, emphasizing the need to preserve these unique habitats. The historical significance of medicinal plants in ancient civilizations and traditional healing systems like Ayurveda underscores their relevance in contemporary healthcare. The study emphasizes the urgent need to conserve these natural resources due to urbanization, deforestation, and climate change. The study also highlights the therapeutic potential of medicinal plants and the



importance of interdisciplinary collaboration and community engagement in conservation efforts. The research calls for concerted action to protect the ecological richness of the Shivalik Hills and preserve their medicinal heritage.

REFERENCES

1. Adhikari, P.P. and Paul, S.B. (2018) 'History of Indian Traditional Medicine: A Medical Inheritance.', Asian Journal of Pharmaceutical and Clinical Research, 11(1), p. 421.
2. Bhandari, B. S., & Bijlwan, K. (2022). Vulnerability of forest vegetation due to anthropogenic disturbances in Western Himalayan Region of India. In Research Anthology on Ecosystem Conservation and Preserving Biodiversity (pp. 1297-1312). IGI Global.
3. Bhandari, M. S., Meena, R. K., Shankhwar, R., Shekhar, C., Saxena, J., Kant, R., ... & Ginwal, H. S. (2020). Prediction mapping through maxent modeling paves the way for the conservation of Rhododendron arboreum in Uttarakhand Himalayas. Journal of the Indian Society of Remote Sensing, 48, 411-422.
4. Bhatta, S. P., & Devkota, A. (2020). Community structure and regeneration status of Sal (Shorea robusta Gaertn.) forests of Dadeldhura district, Western Nepal. Community Ecology, 21(2), 191-201.
5. Dhakad, A. K., & Chandra, A. (2018). Impact of Lantana camara Linn. on plant diversity in Shiwalik hills of Haryana. International Journal of Environmental Science, 8(2), 199-204.
6. Haq, S. M., Khan, I., Malik, Z. A., & Singh, B. (2022). Plant diversity and species distribution pattern across the Pir Panjal Mountain forest range in the Western Himalayas. In Biodiversity, Conservation and Sustainability in Asia: Volume 2: Prospects and Challenges in South and Middle Asia (pp. 67-84). Cham: Springer International Publishing.
7. Jha, S. K., Negi, A. K., Alatalo, J. M., Prabhu, V., Jha, M. B., & Kumar, H. (2022). Forest degradation index: A tool for forest vulnerability assessment in Indian Western Himalaya. Sustainability, 14(23), 15655.
8. Kumar, A., Patil, M., Kumar, P., Kumar, M., & Singh, A. N. (2022). Plant ecology in Indian Siwalik range: a systematic map and its bibliometric analysis. Tropical Ecology, 63(3), 338-350.
9. Kumar, R., Verma, R. K., Kumari, K., Kumar, V., & Prakash, R. (2020). Impact of Burning in Winter on Diversity of Ground Flora after Monsoon Period in Mid Altitude Chirpine Zone of Nahan Circle of Himachal Pradesh. Int. J. Adv. Res. Biol. Sci. 2020c, 7(11), 1-11.
10. Pandey, M. M., Rastogi, S., & Rawat, A. K. S. (2013) Indian Traditional Ayurvedic System of Medicine and Nutritional Supplementation. Evidence-Based Complementary and Alternative Medicine, 2013, 1-12.
11. Raju, S. K., Kumar, S., Sekar, P., Murugesan, M., Karthikeyan, M., Elampulakkadu, A., & Arthanari, M. (2022) Therapeutic and Pharmacological Efficacy of Achyranthes aspera Linn.: An Updated Review. Journal of Drug Delivery and Therapeutics, 12(3), 202–214.
12. Schmerbeck, J., & Gupta, S. (2022). Lantana camara does not exclude all tree regeneration: An opportunity to help establish mixed forests in the Western Himalaya?. Forest Ecology and Management, 520, 120366.
13. Shah, S. (2019) Ayurveda: The Conventional Indian Medicine System and its Global practice. International Journal of Innovative Science & Technology, 4(1), 13–33.
14. Singh, S. (2019). Forest landscape characterization for biodiversity conservation planning and management gaps in Northwestern himalaya using geospatial technology. Remote sensing of Northwest Himalayan ecosystems, 197-236.
15. Singh, S., Singh, B., Surmal, O., Bhat, M. N., Singh, B., & Musarella, C. M. (2021). Fragmented forest patches in the Indian Himalayas preserve unique components of biodiversity: investigation of the floristic composition and phytoclimate of the unexplored Bani Valley. Sustainability, 13(11), 6063.